

- DETHIER, V. G. 1937. Cannibalism among lepidopterous larvae. *Psyche* 44: 110–115.
- DOWNEY, J. C. 1962. Myrmecophily in *Plebejus (Icaricia) icarioides* (Lepid.: Lycaenidae). *Ent. News* 73(3): 57–66.
- D . . . S. 1785. Entomologische Fragmente. N. Mag. Liebh. Ent. 2: 383–384.
- EHRHARDT, R. 1914. Über die Biologie und Histologie der myrmekophilen Organe von *Lycaena orion*. *Ber. Naturf. Ges. Freiburg, i. Br.*, 20: XC–XCVIII.
- FIORI, G. 1958. "Strymon ilicis" Esp. (Lepid. Lycaenidae). *Boll. Ist. Ent. Univ. Bologna* 22(1957): 205–256.
- GUENÉE, M. 1867. D'un organe particulier que présente une chenille de *Lycaena*. *Ann. Soc. Ent. Fr.* (4) 7: 665–668.
- HINTON, H. E. 1951. Myrmecophilous Lycaenidae and other Lepidoptera—a summary. *Proc. South Lond. Ent. Nat. Hist. Soc.* (1949–50) pp. 111–175.
- IWASE, T. 1953. Aberrant feeders among Japanese Lycaenid larvae. *Lepid. News* 7: 45–46.
- 1955. The sixth aberrant feeder in Japan—*Spindasis takanonis* (Lycaenidae). *Lepid. News* 9: 13–14.
- LAMBORN, W. A. 1914. On the relationship between certain West African insects, especially ants, Lycaenidae and Homoptera. *Trans. Ent. Soc. Lond.* (1913) pp. 436–498.
- LENZ, F. 1917. Der Erhaltungsgrund der Myrmekophilie. *Z. indukt. Abst. Vererbungsl.* 18: 44–46.
- MALICKY, H. 1969. Versuch einer Analyse der ökologischen Beziehungen zwischen Lycaeniden (Lepidoptera) und Formiciden (Hymenoptera). *In preparation*.
- NEWCOMER, E. J. 1912. Some observations on the relations of ants and lycaenid caterpillars, and a description of the relational organs of the latter. *Jour. New York Ent. Soc.* 20: 31–36.
- ROSS, G. N. 1964. Life history studies on Mexican Butterflies. II. Early stages of *Anatole rossi*, a new myrmecophilous metalmark. *Jour. Res. Lepid.* 3: 81–94.
- THOMANN, H. 1901. Schmetterlinge und Ameisen. Beobachtungen über eine Symbiose zwischen *Lycaena argus* L. und *Formica cinerea* Mayr. *Jb. Naturf. Ges. Graubunden* 44: 1–40.

OCCURRENCE OF COENONYMPHA INORNATA (SATYRIDAE) IN MAINE

It has been noted in recent years that *Coenonympha inornata* (Edwards) has been expanding its range in the northeastern portion of the United States and Canada.

On July 5, 1968 I collected a female of this species in a roadside bog between Enfield and Passadumkeag, Penobscot Co., Maine. The specimen, which was taken in the early afternoon, appeared to be about one to two days old. Poor weather conditions and a subsequent collecting trip to Canada prevented further collecting in this area.

The specimen has been determined by F. Martin Brown as being closest to *C. inornata inornata* (Edwards). As far as can be determined, this is the first record for this species in the State of Maine.

CLIFFORD D. FERRIS, *Laramie, Wyoming.*